



EVALUATION OF SERUM LEVELS OF LIPID PROFILE IN PATIENT WITH CHRONIC PERIODONTITIS PATIENTS

Zainab J. Raheem

Department of Periodontics, College of Dentistry, Baghdad University.

ABSTRACT

Chronic Periodontitis, a prevalent oral disease, chronic infections has been linked to several systemic health changes, like changing in lipid metabolism. Aims of the study were to evaluate the serum levels of lipid profile (total cholesterol, triglycerides high density lipoproteins, low density lipoproteins and very low density lipoproteins). In study of control groups and compare between them. Also, test the correlation between the serum levels of lipid profile with clinical periodontal parameters. 50 subjects, males and females were included in this study with age range (35-50) years old, they were divided into two groups [CP group: chronic periodontitis group (25 patients)] and control group: (25 systemically healthy subjects, have healthy periodontium). Periodontal health status was determined by measuring the following clinical periodontal parameters (Plaque index (PLI), Gingival index (GI), Bleeding on probing (BOP), Probing pocket depth (PPD) and Clinical attachment level (CAL)). Serum levels of lipid profile were measured by enzymatic colorimetric methods. The results showed that mean values of (TC, TGL, HDL, LDL and VLDL) were high in CP group than control group with highly significant difference at $P < 0.001$. there were TC showed positive, strong and significant with (PLI, PPD and CAL), while with (GL and BOP score 1) showed positive but non-significant correlation. The serum levels of TGL showed positive weak non-significant with (PLI, BOPscore1, and PPD, CAL) and negative correlation with GI. HDL, LDL and VLDL showed positive but non-significant correlation with clinical periodontal parameters except the correlation between LDL and PPD was positive, strong and significant. The elevated levels of serum lipid profile which were observed in chronic periodontitis patients suggest an increased risk for cardiovascular disease in chronic periodontitis patients.

KEYWORD: Chronic periodontitis, serum lipid level, hyperlipidaemia, cholesterol, triglyceride, HDL, LDL.

INTRODUCTION

Chronic periodontitis is a bacterial infectious disease that affects the supporting structures of teeth leading to transient and recurrent bacteremia which lead to many local and systemic inflammatory responses. Periodontitis is a common oral disease in adults that affect many systemic health changes; involving altering lipid metabolism^[1]. Lipids may modify immune response, by altering the macrophages cell membrane, interfering with enzyme systems and proinflammatory cytokines, such as tumor necrosis factor-alpha (TNF- α) and interleukin (IL)-1b^[2], which are associated with periodontal disease^[3]. Many studies revealed an association between periodontitis and hyperlipidemia. Some studies^[4-7] report that patients with hyperlipidemia (diagnosed by physician had higher periodontal parameters compared to systemically healthy controls. The purpose of this study was to show the relation between chronic periodontitis and serum level of total cholesterol (TC), triglycerides (TGL) high density lipoproteins (HDL), low density lipoproteins (LDL) and very low density lipoproteins (VLDL).

MATERIALS & METHODS

In these study Fifty (50) subjects, males and females, aged (35-55) years old were recruited in this study. They were from attendants patients were seeking periodontal treatment in department of periodontics at Teaching Hospital College of Dentistry, Baghdad University. All the participants were informed about the aim of these

Investigations and consented to its protocol. The subjects were divided into: CP group consists of thirty patients diagnosed to have chronic periodontitis and didn't have history of any systemic diseases. Chronic periodontitis in patients was defined as the presence of at least four sites with PPD of (4) mm and clinical attachment loss of (1-2) mm, this made according to the international classification system for PD^[8]. Control group consists of twenty patients with clinically healthy Periodontium this was defined by GI scores < 0.5 ^[9] with no pockets or clinical attachment loss and no history of any systemic diseases. The body mass index of all 50 participants was assessed depending on the criteria by Nieto Garcia *et al.*^[10]. Any participants had BMI more than 25 excluded from the study; also the other exclusion criteria were included history of other chronic, systemic diseases with known associations with PD as diabetes mellitus, Rheumatoid Arthritis *etc.* Smoker medication (anti-inflammatory or antimicrobial therapy) within previous 3 months, periodontal treatment within previous 3 months. Pregnancy, Contraceptive pills. Clinical periodontal parameters examination was done by using Michigan O periodontal probe on four surfaces (mesial, buccal/ labial, distal and lingual/ palatal) of all teeth except third molar, all subjects must have at least 20 teeth. The collected data include assessment of (PLI)^[11] (GI)^[9] BOP^[12] PPD^[8] CAL^[8]. After the clinical periodontal parameters examination 5ml venous blood was collected from study and control groups after a minimum twelve-hour fast, for

Lipid profile of patient with chronic periodontitis patients

the biochemistry analysis of the serum lipid profile levels {TC, TGL, HD, LDL}. Enzymatic colorimetric method using (AGAPPE, Indea) kit. VLDL was assessed

according to the formula in which VLDL=TC/5). Normal values were taken as follows.

Enzyme	Normal range
TC	150 – 220 mg/dL
TGL	Male : 60-165 mg/dL Female : 40-140 mg/dL
LDL	Male : 35 - 80 mg/dL Female: 42 - 88 mg/dL
HDL	Desirable < 130 mg/dL Borderline High Risk for CHD 130-159 mg/dL High Risk for CHD >160 mg/dL.
VLDL	2-30 mg\dl

Statistical analysis was assessed using Mean, standard deviation (SD), Mean of percentage T-test, and Pearson coefficient of correlation).

RESULTS

Table (1) showed Descriptive statistics of the clinical periodontal parameters (PLI, GI, PPD, CAL and BOP score 1) for study (CP) and control groups. This study showed that serum levels of lipid profile were higher in chronic periodontitis patients than control group with highly significant difference between the groups as demonstrated in table (2). The serum levels of TC was (219.48± 31.02), TGL (113.85±11.87), HDL(45.36±6.23), LDL(152.25 ±9.61), VLDL(30.45 ±1.92) in CP group while in Control

group were {(155.32±12.21), (89.03±15.58), (42.41 ±1.49), (76.25 ±13.82), (15.25 ±2.76)} respectively. Table (3) revealed the correlation between the serum levels of lipid profile and clinical periodontal parameters in CP group, serum level of TC showed positive, strong and significant with (PLI, PPD and CAL) ,while with (GL and BOP score 1) showed positive but non-significant correlation. The serum levels of TGL showed positive weak non-significant with (PLI, BOPscore1, PPD, CAL) and negative correlation with GI. HDL, LDL and VLDL showed positive but non-significant correlation with clinical periodontal parameters except the correlation between LDL and PPD was positive, strong and significant.

TABLE 1: Descriptive statistics of the clinical periodontal parameters for study groups

Groups	PII		GI		PPD		CAL		BOP
	Mean	+ SD	Percentage Score 1						
CP	1.73	±0.6	1.81	±0.35	5.06	±0.91	3.09	±0.98	53.72%
Control	0.8	±0.09	0.74	±0.15					

TABLE 2: Descriptive statistics of the serum levels of lipid profile for study groups with Comparison between the groups

Variable	CP group		Control group		t-test	P-value	Sig.
	Mean	+ SD	Mean	+ SD			
TGL (mg\dl)	219.48	±31.02	155.32	±12.21	9.63	<0.001	HS
TGL(mg\dl)	113.85	±11.87	89.03	±15.58	6.337	<0.001	HS
HDL(mg\dl)	45.36	±6.23	42.41	±1.49	22.56	<0.001	HS
LDL(mg\dl)	152.25	±9.61	76.25	±13.82	2.3	0.02	S
VLDL(mg\dl)	30.45	±1.92	15.25	±2.76	6.34	<0.001	HS

TABLE 3: Pearson correlation between serum levels of lipid profile and clinical periodontal parameters at study group

Variables		PLI	GI	PPD	BOPI	CAL
TC	r	0.564	0.209	0.729	0.057	0.915
	P value	0.003	0.316	<0.001	0.78	<0.001
	Sig.	S	NS	HS	NS	HS
TGL	r	0.235	-0.173	0.0185	0.333	0.3
	P value	0.258	0.408	0.93	0.1	0.145
	Sig.	NS	NS	NS	NS	NS
HDL	r	0.0831	0.142	0.0511	-0.12	0.374
	P value	0.69	0.498	0.808	0.57	0.065
	Sig.	NS	NS	NS	NS	NS
LDL	r	0.292	0.0166	0.398	0.004	0.1537
	P value	0.156	0.937	0.048	0.98	0.462
	Sig.	NS	NS	S	NS	NS
VLDL	r	-0.17	0.0188	-0.254	0.004	0.0577
	P value	0.416	0.928	0.22	0.98	0.784
	Sig.	NS	NS	NS	NS	NS

DISCUSSION

The present study demonstrated higher serum levels of lipid profile in CP group than in healthy subjects. Many studies address a relation between chronic periodontitis and serum levels of lipid profile^[13-19] and found elevated serum levels of lipid profile in CP. Each of these studies has their own specifications such as definition of periodontal disease definition, study design and the assay for serum lipid levels assessment. Periodontal disease is the second most common oral disease that is initiated by specific species of pathogenic microorganisms in the dental biofilm. Inflammation is the central cause for tissue destruction that occurs in chronic periodontitis. In the periodontal tissues, the inflammatory process starts with the penetration of the various bacterial toxins and enzymes through the lining epithelium. The inflammatory process becomes chronic when the pathogenic microorganisms continue to grow and cannot be eliminated by the host response. Thus, the inflammatory reaction persists leading to damage to the periodontal tissues. Chronic periodontitis is thus characterized by persistent gingival inflammation, fibrosis, and loss of periodontal tissues. If not treated, it results in apical migration of the functional epithelium and destruction of the periodontal attachment apparatus including the alveolar bone^[20-21]. The two possible mechanisms suggested by which this can occur are by direct migration and colonization of the pathogenic microorganisms to distant organs, leading to an inflammatory reaction at sites distant from the point of infection, and also the systemic inflammation as a result of metastatic periodontal inflammation^[22]. Thus, the present study was conducted to evaluate the lipid profile (TC, TGL, HDL, LDL and VLDL) of the groups to understand the association between chronic periodontitis and dyslipidemia.

Iacopino *et al.* showed any local inflammation causes raised in systemic pro-inflammatory levels will be a potential factor of lipid imbalance, as in chronic periodontitis causes rising in systemic levels of the IL-1 and TNF- which promote the lever to biosynthesis of cholesterol^[21]. Fentoglu *et al.* established the accuracy of this fact by demonstrating the association between IL-1, IL-6 and TNF- levels and TC/HDL ratio^[23]. On the other hand, the periodontal pathogens and their systemic by-products have direct effect by elicitation an immune-inflammatory response and changes in the lipid metabolism, leading to higher levels of VLDL and LDL and decrease in the level of HDL increasing LDL oxidation and the expression of receptors for LDL involved in the control of plasma lipid levels^[24, 25]. In periodontitis there were increased in production of ROS that would contribute to an increase in the process of lipid peroxidation. These Different mechanisms have been proposed above could be explained the elevated serum levels of lipid profile in CP group. As chronic periodontitis its chronic in nature with local and systemic immunologic host responses, there were sustained rise in systemic levels of the pro-inflammatory cytokines that would be led to abnormal serum lipid levels and that might explain the positive correlation between serum levels of lipid profile and clinical periodontal parameters as with sever periodontal destruction increased in systemic levels

of inflammatory cytokines which in return effect on lipid peroxidation. In this study the exact mechanism of association between chronic periodontitis and elevated serum levels of lipid profile was not clarified and it needs to be further explored, beside that the etiology of hyperlipidaemia is multifactorial, therefore, several factors could affect the result, specially high serum lipid profile levels as an inappropriate dietary regimen being included in the study. Although, in our study, overweight and obese participants were excluded this might minimize the possibility of participants with an inappropriate dietary regimen being enrolled in the study.

REFERENCES

- [1]. Ana Cristina Posch Machado, Maria Rozeli de Souza Quirino, Luiz Fernando Costa Nascimento. Relation between chronic periodontal disease and plasmatic levels of triglycerides, total cholesterol and fractions. *Braz Oral Res* 2005; 19(4):284-9.
- [2]. Cutler, C.W., Shinedling, E.A., Nunn, M., Jotwani, R., Kim, B.O., Nares, S. (1999) Association between periodontitis and hiperlipidemia: cause or effect? *J Periodontol.* 70:1429-34.
- [3]. Losche, W., Karapetow, F., Pohl, A., Pohl, C., Kocher, T. (2000) Plasma lipid and blood glucose levels in patients with destructive periodontal disease. *J Clin Periodontol*; 27:537-41.
- [4]. Noack, B., Jachmann, I., Roscher, S. (2000) Metabolic diseases and their possible link to risk indicators of periodontitis. *J Periodontol* 2000; 71:898-903.
- [5]. Awartani, F., Atassi, F. (2010) Evaluation of periodontal status subjects with hyperlipidemia. *J Contemp Dent Pract.*, 11:33-40.
- [6]. Chu, X., Newman, J., Park, B., Nares, S., Ordonez, G., Iacopino, A.M. (1999) In vitro alteration of macrophage phenotype and functions by serum lipids. *Cell Tissue Res.*, 296:331-337.
- [7]. Heasman, P.A., Collins, J.G., Offenbacher, S. (1993) Changes In crevicular fluid levels of interleukin-1beta, leukotriene B4, prostaglandin E2, thromboxane B2, and tumour necrosis factor alpha in experimental gingivitis in humans. *J Periodontal Res.*; 28:241-247.
- [8]. American Academy of Periodontology (AAP). Severity of chronic periodontitis. *Annals of Periodontology*, 1999; page 38.
- [9]. Loe, H. (1967) The Gingival Index, the Plaque Index and the Retention Index Systems. *Journal of Periodontology*, 1967; 38(6):610-616.
- [10]. Nieto-García, F.J., Bush, T.L., Keyl, P.M. (1990) Body mass definitions of obesity: sensitivity and specificity using self-reported weight and height, 1(2): 146-52.
- [11]. Silness, J., Loe, H. (1964) Correlation between oral hygiene and periodontal condition. *Acta OdontolSc and*, 1964; 22: 121-135.
- [12]. Newbrun, E. (1996) Indices to measure gingival bleeding. *Journal Periodontology*, 1996; 67(6):555-561.
- [13]. Lee, J.B., Yi, H.Y., Bae, K.H. (2013) The association between periodontitis and dyslipidemia

- based on the Fourth Korea National Health and Nutrition Examination Survey. *J Clin Periodontol.*, 40:437–42. [PubMed]
- [14]. Deepu Mathews Panickal, Prashanth Evuru, Jeethu John Jerry, Little Mahendra, Sandhya Cherukuri, Jaideep Mahendra IJCRR. Estimation of serum lipid levels in chronic periodontitis patients: A case-control study. 2016; 8(8): 23-26.
- [15]. Ferial Taleghani, Mahmoud Shamaei, Masoud Shamaei *Acta Medica Iranica Association between Chronic Periodontitis and Serum Lipid Levels*, 2010. 48(1):47-50.
- [16]. Saravanan, V., Ravishankar, P.L., Pradeep Kumar, K. Rajapandian, V. Kalaivani, and M. Prem Blaisie Rajula-Estimation of Serum Triglycerides, Serum Cholesterol, Total Protein, IgG Levels in Chronic Periodontitis Affected Elderly Patients: A Cross-Sectional Study. *J Int. Soc. Prev. Community Dent.* 2017 Mar-Apr; 7(2): 120–124.
- [17]. Lal, V., Dubey, D., Rath, S.K., Lohra, P. (2015) Effect of chronic periodontal infection on systemic lipid profile: A clinical and biochemical study. *J Int Clin Dent Res Organ* 2015; 7:106-10.
- [18]. Sandi, R.M., Pol, K.G., Basavaraj, P. Nitin Khuller, and Shilpi Singh *J Clin Diagn Res.* (2014) Association of Serum Cholesterol, Triglyceride, High and Low Density Lipoprotein (HDL and LDL) Levels in Chronic Periodontitis Subjects with Risk for Cardiovascular Disease (CVD): 8(1): 214–216.
- [19]. Hamissi, J., Shahsavarani, M.T., Shahsavarani, H., Sayahpour, S., Hamissi, H.A. (2010) comparison of serum lipid level between patients with periodontitis and healthy individual. *Periodontol Implant Dent.*, 2:29–32.
- [20]. Biju Thomas, Rajendra, B. Prasad, Sucheta Shetty, and Vishakh, R. (2017) Comparative Evaluation of the Lipid Profile in the Serum of Patients with Type II Diabetes Mellitus and Healthy Individuals with *Contemp Clin Dent.* 8(1): 96–101. 101.
- [21]. Iacopino, A.M., Cutler, C.W. (2000) Pathophysiology relationships between periodontitis and systemic diseases: Recent concepts involving serum lipids. *J Periodontal.* 2000; 71:1375–84. [Pub Med].
- [22]. Hayashi, C., Gudino, C.V., Gibson, F.C., 3rd, Genco CA. Review: Pathogen-induced inflammation at sites distant from oral infection: Bacterial persistence and induction of cell-specific innate immune inflammatory pathways. *Mol Oral Microbiol.* 2010; 25:305–16.
- [23]. Fentoglu, O., Bozkurt, F.Y. (2008) The bi-directional relationship between periodontal disease and hyperlipidemia. *Eur J Dent* 2008; 2: 142 – 146.
- [24]. Maekawa, T., Takahashi, N., Tabeta, K. Chronic oral infection with *Porphyromonas gingivalis* accelerates atheroma formation by shifting the lipid profile. *PLoS ONE* 6(5), e20240 (2011).
- [25]. Fentoglu, O., Koroglu, B.K., Hicyilmaz, H. (2011) Proinflammatory cytokine levels in association between periodontal disease and hyperlipidaemia. *J. Clin. Periodontol.* 38(1), 8–16.